




Founder effects determine the genetic structure of the water flea *Daphnia* in Ethiopian reservoirs

Tsegazeabe H. Haileselasie ^{1,2} Joachim Mergeay ^{1,3*} Joost Vanoverbeke,¹ Luisa Orsini ⁴,
Luc De Meester¹

¹Laboratory of Aquatic Ecology, Evolution and Conservation, KU Leuven, Leuven, Belgium

²Department of Biology, College of Natural and Computational Science (CNCS), Mekelle University, Mekelle, Ethiopia

³Research Institute for Nature and Forest, Geraardsbergen, Belgium

⁴School of Biosciences, University of Birmingham, Edgbaston, Birmingham, UK

Abstract

Founder effects introduce stochasticity in the genetic structure of species at the regional scale. To the extent that founder effects are important, they will result in a reduced signature of space, time, and environmental variation in landscape genetic data. We studied the metapopulation genetic structure of recently founded populations of the microcrustacean *Daphnia sinensis* in 10 Ethiopian water reservoirs. We used three different approaches to estimate the number of effective founders and applied them to two independent genetic marker sets in an effort to investigate the role of founder effects and to estimate effective size of the founding population. Estimates of founding sizes rarely exceeded eight individuals but were most often limited to less than four individuals. No associations of genetic identities, gene frequencies, measures of genetic diversity, or differentiation with environmental and spatial variables were found. Age and size of the reservoirs were not correlated with genetic diversity measures or number of founders in these reservoirs. These findings indicate that neither strong selection, nor dispersal limitation are responsible for the observed pattern of genetic variation. Our results suggest a regional population structure that is strongly impacted by founder events, reflecting colonization by just a few founders per waterbody, and not noticeably influenced by subsequent dispersal and gene flow. Our results show that rapid colonization of empty habitats and fast population growth by a handful of founders can result in strong founder effects, even in relatively large habitats that are likely regularly reached by new immigrants.

Metapopulation theory describes the interplay between colonization and extinction rates on patch occupancy as a function of death, birth, and dispersal rates (Levins 1969; Hanski 1998). Population genetics, on the other hand, considers occupied patches and how gene flow and population size interact to influence genetic structure within and among demes of a metapopulation (Wright 1949). In reality, both extinction-colonization dynamics of local populations and changes in genetic diversity by gene flow and drift within these local populations act simultaneously. Dispersal, which is one of the most fundamental processes in ecology, affects many aspects of evolution and population genetics if translated into successful colonization (Bilton et al. 2001). Dispersal allows individuals to establish new populations in an

empty patch and promotes range expansion following colonization of new sites.

In a metapopulation genetics context, the colonization of an empty habitat patch by founders can be considered as a special case of gene flow. As the new local population grows in size to carrying capacity, new neutral immigrants (having equal expected fitness) can still enter the population, but their relative contribution to the local gene pool is expected to decrease as local population size at the time of immigration becomes larger. In this initial colonization scenario, at least two processes are responsible for successful colonization and establishment of a population in an empty patch. First, the response of the immigrants to the local environmental conditions of the habitat they colonize. Second, differences in their time of arrival at the site, which generates a numerical advantage to the first colonizers over late-comers (Boileau et al. 1992; De Meester et al. 2002). In contrast to the relative ease of establishment of founders, which hardly experience intraspecific competition, individuals attempting

*Correspondence: joachim.mergeay@inbo.be

Additional Supporting Information may be found in the online version of this article.

to immigrate into an established population close to carrying capacity are faced with strong intraspecific competition by the resident population and low levels of resources. Due to this, realized rates of gene flow may be much lower than expected on the basis of the rates of dispersal (De Meester et al. 2002). This reduced establishment success may strongly contribute to prolonged persistence of founder effects (Boileau et al. 1992; De Meester et al. 2002; Ventura et al. 2014). Because founder effects represent a type of sampling error, for the mere reason that the source (local or regional) population is sampled by a very low number of individuals to start a new population, they introduce stochasticity in the genetic structure at the regional scale, which tends to result in a reduced signature of space and environmental variation in the genetic data (Orsini et al. 2013).

Estimates of the number of founders represent baseline estimates for ecological dispersal rates (m_c , the observed number of migrants), which represent the maximal potential for gene flow (m_e , the effective number of migrants) among populations. Gene flow, the realized effect of ecological dispersal on genetic structure, can be estimated indirectly through population genetics as well (Broquet and Petit 2009). Although we know from many population genetic studies in zooplankton that gene flow (m_e) is often much lower than expected, we have relatively few good estimates of ecological dispersal rates (m_c), because they are so hard to measure directly (Bilton et al. 2001). Nevertheless, good dispersal estimates provide baseline information for a broad array of ecological and evolutionary studies (Broquet and Petit 2009). Distinguishing between dispersal and gene flow is essential, especially for biological conservation of populations and species.

Here, we take advantage of the recent creation of water reservoirs in Northern Ethiopia and the colonization of these water bodies by zooplankton to estimate the number of founders of populations of a zooplankton species using genetic methods. The reservoirs studied here are young (6–18 yr) and two to three orders of magnitude larger than most other systems studied so far on founder effects in zooplankton and small invertebrates (Boileau et al. 1992; Haag et al. 2006; Louette and De Meester 2007). Specifically, we present patterns of genetic composition and differentiation of the water flea *Daphnia sinensis* in reservoirs that range in size from 1.8 ha to 45.4 ha. Using variation at nuclear (nDNA) and mitochondrial (mtDNA) genetic markers, we estimate allele frequencies, within-population genetic variation and among-population genetic differentiation, and relate genetic variation and genotype composition to spatial, environmental, and temporal variables. We use various methods to independently estimate founding population sizes, and thereby provide baseline estimates of dispersal rates. Using information on the observed genetic structure (F_{ST}) and the associated expected gene flow at various levels of migration-drift equilibrium and ages of populations, we estimated the number of effective founders.

Methods

Study region and sampling

The studied reservoirs are part of a set of reservoirs constructed between 1984 and 2001 in the highlands of Tigray Regional State, Northern Ethiopia. The rainfall in Tigray region is highly seasonal (2 months of heavy rains per year, 10 months of drought) and erratic (because of large year-to-year variation), resulting in moisture stress that hampers the rain-fed agriculture (Haregeweyn et al. 2006). Agricultural development through irrigation has therefore been a priority for the Regional Government of Tigray. The construction of reservoirs was initiated with the main aim to bring food self-sufficiency to the area through irrigation, but the reservoirs have additional functions such as watering cattle and the use of water for household purposes (Asmelash et al. 2007). Thirty-two of these reservoirs have been the subject of a detailed limnological survey (Asmelash et al. 2007; Dejenie et al. 2008). Apart from a single natural lake, not inhabited by the focal species of this study, *D. sinensis*, no similar large and deep aquatic systems are known from Tigray (Dejenie et al. 2008). Naturally, this species occurs in temporary pools and ponds as well as larger temporary ponds and lakes (Gu et al. 2013). The rapid colonization of these reservoirs shortly after their creation by a considerable number of zooplankton taxa, including typical lake species (Dejenie et al. 2008), despite a regional lack of similar habitats suggests that dispersal rates are relatively high and long-distance dispersal events are rather frequent. Water birds (members of the Podicipidae, Pelecanidae, Ciconiidae, Anatidae, and Charadriidae families) are common in and alongside the reservoirs (Dejenie et al. 2008) and are probably important vectors of dormant propagules of zooplankton (Figuerola and Green 2002).

Thirty-two reservoirs were sampled for zooplankton, using a Schindler-Patalas plankton trap, in September 2005 (Dejenie et al. 2008). Ten of these samples contained *D. sinensis* in large enough numbers for population genetic analyses (Supporting Information Table S1). In addition, five temporary natural wetlands were sampled, two of which contained *D. sinensis* (henceforth called T1 and T3), bringing the total number of independent samples to 12. All *Daphnia* samples were preserved in 100% ethanol until further processing. Although previously identified as *D. carinata* King by Dejenie et al. (2008), DNA barcoding indicates that individuals from these reservoirs belong to *D. sinensis*, a member of the *Daphnia similis* species complex (Popova et al. 2016). Measurements of geographic position and morphometric, physical, chemical, and biotic variables were recorded for each sampled reservoir (Supporting Information Table S1; Dejenie et al. 2008). Age of the reservoirs was expressed as number of years at sampling time since construction of the reservoir. The two natural populations T1 and T3 were first excluded from all age related analysis, and were in a second analysis arbitrarily given the same age as the oldest reservoir.

Table 1. Genetic diversity in mtDNA haplotypes: observed frequencies (expressed as fractions) of each haplotype per waterbody and diversity descriptors. Average observed alpha diversity is the average observed within-sample diversity weighted by sample size. Average expected alpha diversity gives the expected value of HR or HD given a panmictic population over all waterbodies, using 10,000 permutations. The range of the expected values shows the lowest and highest value among all permutations over individuals. All observed values deviate significantly ($p < 0.0001$) from expected values. True beta diversity is calculated as gamma/alpha.

Waterbody	N	Haplotype number						HR	HD-Si
		1	2	3	4	5	6		
Adi Gela	17	0	0.71	0.06	0.18	0.06	0	4	1.86
Adi Kenafiz	18	0	0	0.39	0.56	0	0.06	3	2.16
Dibla	27	0.26	0.07	0.67	0	0	0	3	1.93
Gereb Awso	35	1.00	0	0	0	0	0	1	1.00
Gereb Mihiz	31	0.45	0.16	0.39	0	0	0	3	2.63
Gum Selasa	33	0.27	0.06	0.27	0.39	0	0	4	3.25
Haiba	6	0.83	0	0.17	0	0	0	2	1.38
Mai Leba	29	0.03	0.93	0	0.03	0	0	3	1.15
Meala	32	0.97	0.03	0	0	0	0	2	1.06
Tsinkanet	12	0	1.00	0	0	0	0	1	1.00
Temp 1 (T1)	22	0.14	0.36	0.50	0	0	0	3	2.49
Temp 3 (T3)	23	0	1.00	0	0	0	0	1	1.00
Overall frequency		0.33	0.36	0.203	0.097	0.005	0.005	—	—
Average observed (alpha)		0.38	0.30	0.19	0.12	0.01	0.01	2.5	1.74
Total diversity (gamma)		—	—	—	—	—	—	6	3.43
Average expected alpha		—	—	—	—	—	—	4	3.40
Range expected alpha		—	—	—	—	—	—	3.7–4.2	3.01–3.67
True beta diversity		—	—	—	—	—	—	2.4	2.10

HD-Si, true haplotype diversity measured with the Simpson index; HR, haplotype richness; N, number of individuals extracted per sampling site.

Genotyping

DNA of individual *Daphnia* was extracted using the HotShot protocol (Montero-Pau et al. 2008). Sample sizes ranged from 6 to 35 individuals per population for mtDNA (Table 1; 10 out of 12 samples with 17 or more individuals) and from 15 to 34 individuals per population for nDNA (Table 2). Differences in sample sizes between both markers are due to unsuccessful amplification with either approach. A fragment of 341 nucleotides of the mtDNA Cytochrome Oxidase gene, subunit 1 (*COI*) was amplified using primers *SCox1F1* (GGC CCC AGA TAT GGC TTT) and *SCox1R2* (GCT CCA GCT AAT ACT GGT AAA CTT), specifically designed for this study. The polymerase chain reaction mix of 25 μ L contained 2 μ L DNA, 2.75 μ L 10 \times polymerase chain reaction (PCR) buffer (10 mM Tris-HCl; pH 8.3; 50 mM KCl), 0.4 μ M of each primer, 2.2 mM MgCl₂, 0.2 mM of each deoxynucleotide triphosphates (dNTP), and 1 unit Silverstar *Taq* DNA polymerase (Eurogentec[®], Liege Belgium). PCR cycling conditions, PCR product purification, and sequencing followed the methods of Mergeay et al. (2007). The purified fragments were sequenced using 3.2 pmol of *SCox1F1* primer and the ABI Big Dye Terminator Kit. Sequences were aligned and trimmed in Mega 4.1 (Kumar et al. 2008).

Variation at six microsatellite loci using primers originally developed for the related species *Daphnia magna* (*B088*,

B172, *B087*, *S6-38*, *B064*, and *Dma15*; Agostini et al. 2010) was assessed in a single multiplex PCR reaction of 10 μ L consisting of 5 μ L HotStar *Taq* DNA polymerase buffer (Qiagen[®], Hilden Germany), 0.15 μ M, 0.5 μ M, 0.3 μ M, 0.2 μ M, 0.1 μ M, and 0.3 μ M of each primer of locus *Dma15*, *B087*, *B064*, *S6-38*, *B088*, and *B172*, respectively, and 2 μ L of template DNA. PCR consisted of 15' at 95°C followed by 30 cycles of 30" for each step at 95°C, 56°C, and 72°C, and a final elongation step at 60°C for 30'. Polymorphism was assessed on an ABI PRISM[®] 3130 genetic analyzer (Applied Biosystems[®], Foster City, California, U.S.A.), using an internal Liz Gene-scan size standard by means of the Genemapper 4.0 software (Applied Biosystems[®], Foster City, California, U.S.A.).

Population genetic data analysis

Nucleotide diversity of the sequenced *COI* fragments was calculated in DNAsp (Rozas et al. 2003) version 4.5. Because we were not interested in the evolutionary relationships among haplotypes that originated thousands of years prior to the colonization of these reservoirs, no attempts were made to construct haplotype networks or to calculate genetic differentiation among populations based on haplotype identity. The observed haplotype frequencies were primarily used to estimate the number of founders involved in the colonization of each reservoir. We calculated observed haplotype

Table 2. Clonal and genetic diversity based on microsatellite loci (nDNA).

Waterbody	<i>N</i>	<i>n</i>	CR	CD	CR/ <i>n</i>	CD/ <i>n</i>	<i>A</i>	AR	<i>H_o</i>	<i>H_e</i>	HWE*	<i>F_{IS}</i>
AG	30	20	19	18.18	0.95	0.91	20	2.97	0.38	0.54	0.001	0.302
AK	30	20	15	11.11	0.75	0.56	11	1.76	0.39	0.34	0.335	-0.164
DIB	30	27	19	12.79	0.70	0.47	15	2.29	0.41	0.32	0.085	-0.29
GA	32	31	13	8.50	0.42	0.27	14	1.95	0.34	0.27	0.108	-0.242
GM	36	34	19	6.64	0.56	0.20	19	2.46	0.43	0.39	0.001	-0.106
GS	40	32	27	22.26	0.84	0.70	18	2.63	0.38	0.47	0.000	0.19
HA	16	16	16	16.00	1.00	1.00	17	2.62	0.35	0.41	0.001	0.145
ML	29	23	22	21.16	0.96	0.92	13	2.13	0.57	0.47	0.012	-0.19
MA	30	20	3	1.23	0.15	0.06	13	1.85	0.51	0.29	0.000	-0.755
TS	18	15	11	7.76	0.73	0.52	18	2.92	0.39	0.52	0.000	0.258
T1	26	23	15	8.97	0.65	0.39	14	2.10	0.21	0.24	0.272	0.138
T3	32	32	31	30.12	0.97	0.94	14	2.13	0.45	0.47	0.335	0.032

CR/*n* and CD/*n* refers to clonal richness and diversity, respectively, corrected for sample size expressed as proportion of clones to total individuals genotyped.

A, number of alleles; AR, allelic richness; CD, clonal diversity; CR, clonal richness; *F_{IS}*, fixation index between individuals within local populations; *H_e*, expected heterozygosity; *H_o*, observed heterozygosity; *N*, sample size; *n*, number of individuals with complete genotypic information (six loci) on which calculations of CR and CD were based.

* The numbers are the *p*-value from a goodness of fit to Hardy-Weinberg equilibrium (HWE) expectations test, using Fisher's exact test method.

richness (HR) for each sample as well as haplotype diversity (HD). HD was calculated as the true diversity equivalent of the Simpson concentration (Jost 2007). These values were compared to expected HR and HD under the null hypothesis that all reservoirs form one panmictic population, using 10^4 permutations in Partition (Veech and Crist 2009). This yields alpha (local gene diversity) and beta (average differentiation) estimators that are converted to their true diversity equivalents (Jost 2007).

Standard measures of genetic diversity (number of alleles, allelic richness, AR, per locus and per population across all loci, observed heterozygosity, *H_o* and expected heterozygosity, *H_e*) at six microsatellite loci were assessed in R using diveRsim package (Keenan et al. 2013). Identical multilocus genotypes (MLGs), on the basis of the combined information of six microsatellite loci, in a given waterbody were considered to belong to a single clone. Clonal diversity (CD) was expressed as the true diversity equivalent of the Simpson concentration, clonal richness (CR) as the number of MLGs per waterbody. Moreover, relative CR was calculated per sample corrected for sample size expressed as proportion of clones to total individuals genotyped as $R = (G-1)/(N-1)$, where *G* is the number of genotypes and *N* indicates sample size.

The standardized genetic variance among populations (*F_{ST}*) was calculated according to Weir and Cockerham (1984). We used 500 bootstrap pseudoreplicates to estimate 95% confidence intervals of the *F_{ST}* values. Genetic structure was assessed using the unbiased estimators of Nei and Chesser (1983) of the overall gene diversity (*H_T*) and subpopulation gene diversity (*H_S*).

Spatial, environmental, and temporal correlates of genetic differentiation

To investigate the role of spatial and environmental variables separately and to disentangle the unique contribution of each variable matrix to the genetic structure of the studied populations, we used redundancy analyses (RDA, a linear constrained ordination technique, Dray et al. 2006). In a multivariate variation partitioning analysis, the contributions of local environmental predictors (*n* = 16, variables provided in Supporting Information Table S1) and spatial predictors are tested by generating adjusted redundancy statistics (*R_{adj}²*) in an RDA analysis. A significant effect of environment would imply sorting of clones with different traits and niches along environmental gradients.

Under a model of persistent founder effects, we expect genetic structure to be mainly caused by chance events as dispersal is likely not limiting at the investigated spatial scale. Hence, we expect to find at most a weak spatial genetic structure in the data (Orsini et al. 2013). To test this, with the nDNA allele frequency data, we performed a principle coordinates analysis (PCoA) and then used the population loadings of the first six PCoA axes as dependent variables in a distance-based redundancy analysis (db-RDA). In this RDA, we attempted to explain the observed genetic variation as a function of distance-based eigenvector maps (dbMEM) (Dray et al. 2006). This analysis allows to find spatial patterns in the genetic data other than linear ones, making this a more powerful approach with lower type II error rates than Mantel tests (Legendre and Fortin 2010). Five positive dbMEM eigenvectors were retained and were used as explanatory variables in a forward selection procedure.

Although this particular approach has the risk of identifying a false positive spatial signal (*see* Blanchet et al. 2008), the double stop criterion of Blanchet et al. (2008) is very conservative with regard to small datasets. Here, we take a more liberal approach, involving forward selection of spatial variables without prior testing of the overall spatial model, to make sure that any lack of a detectable spatial signal is not due to the use of conservative statistical methods. Besides, a Mantel test was used to test for any spatial trend in the genetic data.

In parallel, we performed a separate RDA relating the mtDNA data to the spatial data (dbMEM). These mtDNA data were Hellinger-transformed to allow the use of linear regression analyses in zero-inflated data (Legendre and Gallagher 2001). Using nonparametric correlation (Spearman's rho) analyses, we related diversity measures (H_e , AR, HR, HD, CR, CD, number of founders at mtDNA, and number of founders at nDNA) to age, depth, and size of the reservoirs. Under a model of persistent founder effects, genetic diversity should not be related to age or size of the reservoir, as late-arriving immigrants are expected to have little impact on the genetic structure compared to the very first founders. If founder effects still persisted at the time of sampling, we expect that old and large populations do not differ in genetic structure from younger and/or smaller populations. Specifically, we tested if a model of population structure including reservoir age, reservoir size, or their interaction explained the genetic structure better than the null model not including age or size using Geste v. 2 (Foll and Gaggiotti 2006). Geste calculates a specific F_{ST} value for each population that represents the population specific contribution to the total genetic differentiation in the metapopulation. Then, the effect of age and/or size on these F_{ST} values is evaluated using generalized linear models. The posterior probability of each model was used to select the model with the highest probability given our data.

Estimating the number of effective founders

We used three different approaches to estimate the number of effective founders, which reflects how many individuals contributed to the observed genetic diversity in each local population. First, we used a general approach based on F -statistics from microsatellite data, using the principle that the inbreeding coefficient among populations just after colonization is $F_{ST} = (2K)^{-1}$, with K the average number of founders per population (Wade and McCauley 1988; Boileau et al. 1992). Confidence intervals (95% CI) for F_{ST} were calculated by bootstrapping over loci (500 replicates). This method provides the average effective size of the founding population. Second, we used a general simulation approach in the programming environment R (R Core Team 2014) to calculate the expected HR and HD under a model of random colonization from a regional gene pool with 1, 2... 10, 15, and 20 founders. Expected HR and HD values were calculated by

randomly sampling 10^4 times the pre-set number of founders in 200 populations from an estimated regional frequency distribution, which was based on the actual haplotype counts over all water bodies, or on presence-absence data for each haplotype per waterbody. For each of the random samples, we calculated the probability that the expected average HR or HD, over the 200 population, was smaller or larger than the values observed in our empirical dataset. The product of these values provides the overall probability that the observed HD or HR is achieved by the corresponding number of founders. The R-script is available as Supporting Information (Supporting Information Table S2), and provides the average census size of the founding population. Third, we used a population-specific approach for which we used the Colonize script (Mergey et al. 2007). This is a standalone command-line tool that calculates the likelihood that a predefined number of founders from a predefined source population established the focal population, given the gene frequency distribution of the source and the sink populations. The number of founders associated to the highest likelihood score provides the best estimate for the founding propagule size of that population. Again, this provides the census size of the founding population, which ignores that some individuals contributed less to the genetic structure of the founding population than other individuals. For each waterbody, we calculated likelihood scores for one to thirty founders, and set both the number of batches and the number of random samples to 500. Ideally, one has multiple putative source populations from which to sample, so as to assign the most likely source population as well (*see* Mergey et al. 2007 for an example). Here, we have no such prior information, and hence, we use the regional gene pool (the average over all our samples) as the overall source population. To increase the overall robustness of this approach, three different prior allele frequency distributions were used. (1) Using the regional frequency of each allele over the pooled data of all investigated water bodies (distribution = Freq). Here, we used the rare allele correction in Colonize to account for extremely rare alleles. (2) Using presence-absence of each allele per population and counting the frequency of occurrence of each allele over all populations (distribution = Rich). This approach gives less weight to alleles that dominate in certain populations but are rare in other populations. (3) Using three abundance classes for the regional allele frequencies (1: frequency < 15%; 2: frequency 15–30%; 3: frequency > 30%). This approach (distribution = Level) gives even more weight to rare alleles. Analyses with Colonize were performed separately for the mtDNA data and for the nDNA data in order to obtain independent estimates for both marker types. Overall, these three approaches yield one overall estimate based on F_{ST} (first method), two overall estimates based on mtDNA haplotype richness and haplotype distributions (second method), and per reservoir three estimates based on mtDNA and three based on nDNA (third method).

Testing assumptions

All of the outlined methods to estimate the number of founders rely on similar assumptions, but vary in their sensitivity to violations thereof. Here, we outline how we tested for violations of the assumptions. First, we assume that genetic drift has not yet strongly affected allele frequency distributions (especially fixation or loss of alleles), given that they were founded at most 6–18 yr before sampling. Second, we assume that all founders are genetically independent.

The main source of genetic drift in cyclical parthenogens like *Daphnia* is clonal selection (reduction in clonal and genetic diversity as a result of selection among clones in the population) (De Meester et al. 2006), which may erode genetic diversity considerably and thereby reduce our estimates of founding population sizes. To assess whether clonal population structure (leading to a similar signal as genetic bottlenecks) has affected our results, we performed a Spearman Rank order correlation between CD and CR vs. the estimated number of founders per population. This was performed for all estimates of the number of founders, which have different sensitivities for common and rare alleles. Significant correlations would reflect that clonal erosion affects our estimates of founding population sizes. Furthermore, because drift reduces richness faster than diversity (Cornuet and Luikart 1996), founding population size estimates based on richness should be lower than those based on diversity indices if genetic drift is really important. Next, genetic drift affects mitochondrial genetic structure stronger than nuclear genetic structure, because the effective population size at mitochondrial genes is smaller (Hamilton 2011). Hence, our founding number estimates should be lower when using mitochondrial data if these were strongly influenced by genetic drift. We used the Student *t*-test to test for differences among all these cases.

Results

Genetic diversity

Among a total of 285 sequences, we found 25 polymorphic positions out of 299 nucleotide positions in the *COI* gene fragment. This resulted in six distinct mitochondrial haplotypes. Two of these haplotypes, H5 and H6, were singletons detected from Adi Gela and Adi Kenafiz, respectively. Four haplotypes were common, with overall frequency of occurrence of 36.0%, 33.0%, 20.3%, and 9.70% for H2, H1, H3, and H4, respectively (Table 1 and Fig. 1). Most populations, including both natural systems, were dominated by one or two haplotypes (average HD = 1.74). Overall, the observed HD or HR in a given population was always significantly lower (X^2 test, $p < 0.0001$) than the expected diversity or richness assuming a panmictic regional metapopulation (Table 1). Pairwise nucleotide diversity among haplotypes ranged from 0.003 to 0.047 (overall nucleotide diversity = 0.021).

For the microsatellite markers, we found an average of 5.3 ± 3.6 (\pm SD) alleles per locus over the whole metapopulation, whereas the mean allelic richness per locus was 2.58 ± 0.5 per population. The number of alleles per locus ranged from 2 to 11, with a total of 32 alleles scored over the six microsatellite loci combined. The total number of alleles observed across all loci per population ranged from 11 to 20, with a mean allelic richness of 2.32 alleles per locus (Table 2). The observed heterozygosity for the 12 relatively young *Daphnia* populations ranged from 0.21 to 0.57 whereas the expected heterozygosity (H_e) ranged from 0.24 to 0.54 per population (Table 2 and Supporting Information Table S3).

In total, we found 183 unique MLGs out of 293 individuals successfully genotyped. The majority of those MLGs (88%) were represented by a single individual, and another 5% of the MLGs ($n = 10$) were represented by only two individuals.

All values of pairwise genetic differentiation (F_{ST}) were significant ($p < 0.05$). Nearby population pairs were not more related to each other than distant pairs (Supporting Information Table S4). The highest pairwise F_{ST} value was observed in the comparison between T1 and Adi Kenafiz ($F_{ST} = 0.585$), while the lowest pairwise F_{ST} value ($F_{ST} = 0.037$) was between Gereb Awso and Dibla (Supporting Information Table S4).

None of the RDA analyses yielded a model with one or more spatial or environmental explanatory variables that could significantly ($p < 0.05$) explain the variation in the genetic data, either in the distribution of the mtDNA haplotypes, or in the allele frequency data of the microsatellite loci. Exclusion of the two natural systems did not affect the general pattern. Mantel tests between pairwise genetic distance (Nei's genetic distance) and geographic distances or environmental distances yielded Mantel correlation coefficients of $r = -0.114$ ($p = 0.658$) and $r = 0.181$ ($p = 0.212$), respectively, thus confirming the absence of any spatial trend in the genetic data (Fig. 2).

Number of founders

Method 1: F_{ST} -based

The overall among-population fixation index (F_{ST}) was 0.237, with 95% CI ranging from $0.180 < F_{ST} < 0.342$. Without T1 and T3, F_{ST} equaled 0.219 (95% CI: $0.169 < F_{ST} < 0.309$). Since $F_{ST} \approx 1/2N$ at colonization, this reflects average effective founding population sizes of 2.3 individuals (95% CI: $1.6 < N_e < 3.0$). Put differently, the average genetic diversity we observed corresponds to a mean effective founding population size of 1.6–3.0 individuals.

Method 2: Comparing observed to expected richness and diversity estimates

We found an average observed HR = 2.5 and an average observed HD = 1.74 (Table 1). When comparing the average observed levels of HR and HD to expected HR and HD, we found that average founding population size estimates smaller than two and larger than eight are improbable at p -value = 0.05 (Supporting Information Table S5). The highest probability




Fig. 1. Geographic location of the sampling sites and mtDNA haplotype frequencies in each population. Major cities are indicated with a star. Inset on the right shows the overall regional frequency of the six encountered haplotypes. AG, Adi Gela; AK, Adi Kenafiz; Di, Dibla; GA, Gereb Awso; GM, Gereb Mihiz; Ha, Haiba; Me, Meala; ML, Mai Leba; T1, temporary pond 1; T3, temporary pond 3; Ts, Tsinkanet.

scores were obtained with 4, 3, 3, and 4 founders for the four types of simulations (Supporting Information Table S5).

Method 3: Population-specific simulations

We used population-specific simulations using mtDNA and nDNA, based on three prior theoretical allele frequency distributions (Freq, Rich, and Level, in descending order of sensitivity to rare alleles). For mtDNA, all three prior allele frequency distributions yielded very comparable estimates (Table 3 and Supporting Information Table S6), with averages ranging between 3.5 and 4 founders ($1 \leq \text{range} \leq 8$). This did not change appreciably when the natural systems were excluded (results not shown). For nDNA, similar average (2.8–5.5) values were found ($2 \leq \text{range} \leq 13$), although the estimate using the Freq. prior distribution was somewhat higher and was positively skewed due to higher estimates for two populations (Table 3 and Supporting Information Table S6).

Confidence in the prior distribution of regional allele frequencies (Freq.) was unacceptably low (highest observed likelihood score < 0.05) in six cases for nDNA and two cases for mtDNA. The prior distribution based on local richness of alleles (Rich) yielded one estimate at nDNA with too low likelihood scores (Table 3 and Supporting Information Table S6). All these estimates indicate that founding population sizes were typically smaller than five individuals, and very rarely exceeded ten individuals.

Effect of size and age of the reservoir

There was no significant relation ($p > 0.05$) between H_e and surface area or log (surface area) of the waterbody ($S = 183.28$, Spearman rank $r = 0.3563$, $p = 0.126$) or with age of the reservoir ($S = 342.35$, Spearman rank $r = -0.197$, $p = 0.730$, Supporting Information Fig. S1). The only significant correlation found in a total of 42 associations tested (age, depth, area, vs.




Fig. 2. Relationship between Nei's genetic distance and geographic distance (panel **A**; testing for an isolation-by-distance and thus for dispersal limitation; $r = -0.114$; $p = 0.662$) and the Euclidean distance for environmental variables (panel **B**; testing for IBE; $r = 0.181$; $p = 0.212$).

AR, H_e , H_o , HR, HD, CR/N, CD/N, F_{IS} , number of founders at mtDNA [Colonize-Rich], and number of founders at nDNA [Colonize-Rich] was between H_e and average depth ($r = -0.79$, $p < 0.001$). However, after Bonferroni correction, this p -value was larger than 0.05. All other correlations were extremely weak and statistically insignificant at $\alpha = 0.05$ (absolute value of $r < 0.15$, uncorrected $p > 0.10$; Supporting Information Fig. S1). Inclusion of size and/or age of the reservoir did not provide a better model (Geste v. 2) for genetic structure than the more parsimonious null model. All of these results support the hypothesis that founder effects are the main drivers of genetic structure and indicate that clonal genetic drift did not markedly influence our estimates of founding population sizes. In addition, none of the tests we did could show a significant difference between founding

population size estimates based on mitochondrial vs. nuclear genetic data, or based on richness vs. diversity estimates (all p values > 0.05). Thus, we have no indication that the assumptions concerning genetic drift were violated (see Supporting Information Table S7 and Supporting Information Fig. S1).

Discussion

Although we included a broad set of environmental and spatial variables, we found no association between genetic differentiation among the *D. sinensis* populations inhabiting artificial reservoirs in Tigray and environmental variation, space or time (age). Both the nuclear and mitochondrial markers that we used are expected to behave neutrally. As such, we expected a stronger signature of space than of

Table 3. Summary results of colonize analyses with three prior allele frequency distributions (Freq, Rich, Level; see main text for explanation), showing the most likely number of founders for each population, based on either mtDNA or nDNA data, for each population and averaged over all populations. SD, standard deviation. Values with asterisk indicate that the likelihood score was too low ($p < 0.05$) to represent a reliable estimate. Non-integer values represent the average of shared highest scores.

Waterbody	Number of founders with highest likelihood score (colonize)					
	mtDNA			nDNA		
	Freq	Rich	Level	Freq	Rich	Level
Adi Gela	4*	4	4	2*	2	2
Adi Kenafiz	5*	8	7	7*	7	5
Dibla	5	4	4	5	3	2
Gereb Awso	1	1	1	5	2	2
Gereb Mihiz	5	6	4	12.5	4	4
Gum Selasa	8	8	7.5	13	5	4
Haiba	2.5	2	2	3*	2	2
Mai Leba	4	3.5	3	5	3	2
Meala	2	2	2	2*	2	2
Tsinkanet	1	1	1	3*	3*	4
T1	5	5	5	3*	2	2
T3	1	1	1	6	3	2
Average	3.63	3.79	3.46	5.54	3.17	2.75
SD	2.17	2.55	2.23	3.71	1.53	1.14

environment. Still, a correlation with environmental variables may result when particular haplotypes would hitchhike with particular genotypes or fixed allele combinations that are favored under certain conditions. Three studies on strictly asexual zooplankton with comparable sample sizes and statistical power found clear environmental and/or spatial structuring in their studies (Aguilera et al. 2007; Pantel et al. 2011, Haileselesie et al. 2016) suggesting that the lack of patterns in our dataset is not merely a consequence of insufficient statistical power. Evolution-mediated priority effects, the key feature of the monopolization hypothesis (De Meester et al. 2002; De Meester et al. 2016), are expected to be less important in asexual taxa than in similar sexual taxa, due to a reduced ability for rapid local adaptation in the asexuals. As a consequence, it is expected that fitness differences among dispersed clones will lead to a match between environmental gradients and landscape genetic structure in asexual taxa, similar to species sorting in communities (Leibold et al. 2004; De Bie et al. 2012). Conversely, if the colonizing propagules of a sexual species harbors sufficient genetic variation to allow local genetic adaptation, the increased fitness of resident populations may reduce establishment success of new immigrants and thus reduce gene flow (De Meester et al. 2016). As a result, the match between

environmental and genetic variation is expected to be less strong in sexual than in asexual species, which is the emergent pattern from our study on a cyclically parthenogenetic *Daphnia* species and contrasts with the three aforementioned studies that focused on obligately parthenogenetic *Daphnia*.

To investigate the number of founders typically involved in the colonization of new moderately-sized freshwater systems (ranging from 1.8 ha to 45.4 ha in size), we used three different approaches that rely on different test statistics with varying prior parameters, applied on two independent sets of genetic markers. All approaches indicated that typically less than five founders per habitat were responsible for the observed pattern of genetic diversity in the studied reservoirs. Irrespective of whether we used estimates based on richness data or more detailed frequency data, we obtained very similar estimates, showing that our results are robust to strong allele frequency changes that may have occurred since colonization. Admittedly, the different approaches we used all rely on similar assumptions, including an absence of genetic drift since colonization, and genetic independence of each founder. The second assumption that the founders are genetically independent from each other may have been violated to some extent. Birds, for example, may disperse more than one dormant stage at the same time from a single source, thereby introducing multiple related propagules. Especially results from mtDNA are expected to be prone to such bias, given the much lower local and regional genetic variation compared to the levels of variation found at nDNA. Estimated numbers of founders for nDNA and mtDNA were, however, very similar.

We have detected high genetic differentiation among population ($F_{ST} = 0.232$) and no isolation by dispersal limitation. This indicates low levels of gene flow among populations. Furthermore, we failed to detect isolation-by-environment (IBE), which rules out the possibility that sorting of genotypes along environmental gradients similar to species sorting in communities (Leibold et al. 2004) might have driven the observed high genetic differentiation among populations. Thus, our results support the idea that colonization dynamics in a newly created metapopulation are strongly affected by founder effects exerted by a limited number of founding genotypes. The founder effects observed here indicate that metapopulation and colonization dynamics in this species resemble a lottery model (Sale 1977). In Sale's (1977) lottery model, individuals compete for a limited number of discrete resources and once a resource is claimed, an individual cannot be usurped from it. The classic lottery model was formulated at the community level and with respect to microsites. However, it here acts at the level of genetic variants of a species, and at the habitat level in a metapopulation. Populations are thus founded by a small number of individuals from a varied array of regional sources. As long as a local genetic variant persists (also if persistence is mediated through dormant stages; Mergeay et al.

2007), the niche space will continue to be occupied by these local variants, thereby pre-empting niche space for immigrants. Several empirical studies focusing on colonization of novel habitats have shown that dispersal rates in zooplankton are high (Jenkins and Buikema 1998; Cáceres and Soluk 2002; Louette and De Meester 2005). The lack of spatial genetic patterns in our dataset also suggests that dispersal per se is not limiting at the spatial scale here studied.

The sole environmental variable that showed a significant but negative correlation with genetic diversity was average lake depth. One may speculate that the negative correlation between depth and H_e reflects a species-specific preference for shallow waters, thereby reducing the likelihood that a colonizing propagule will survive in deep reservoirs. This is indeed expected from an organism that seems to naturally inhabit shallow pools. In that case, however, we would also expect a similar negative relation between depth and number of founders, or other measures of genetic diversity, which was not the case. An alternative explanation is that deeper lakes result in more stable habitat conditions and therefore in populations that survive year-round and are thus less dependent on dormant egg banks for survival. It is well known that more permanent populations in *Daphnia* exhibit lower genetic diversity because of ongoing clonal erosion (Hebert 1987; De Meester et al. 2006).

Earlier studies (Boileau et al. 1992; Haag et al. 2006) already showed that founder events can strongly determine metapopulation structure, but the habitats they studied were very small (less than $< 100 \text{ m}^2$). The systems we study are thousand times larger than the typical size of the small habitats studied earlier, with associated differences in carrying capacity, effective population size, genetic drift, and inbreeding. Although the results shown here should be interpreted with some caution given that the limited number of reservoirs that was inhabited by the studied *Daphnia* species resulted in a reduced statistical power in detecting spatial and environmental patterns, our analyses strongly indicate that zooplankton populations of these new large waterbodies are typically founded by just a handful of individuals. Interestingly, the number of founders in these reservoirs (on average 4–6) is strikingly similar to the range found in ponds with population sizes that are up to a thousand times smaller (Boileau et al. 1992; Louette et al. 2007). Similarly, the local recolonization by *Daphnia barbata* of the 150 km^2 large Kenyan Lake Naivasha happened most likely by no more than nine individuals from an old dormant egg bank (Mergeay et al. 2007).

Inbreeding effective population size (N_e) in populations is a function of the number of founders and is thus generally small in our zooplankton population. It seems that in zooplankton, habitat size per se, at least within given boundaries, may have little influence on the effective population size. Next to the low number of founders that seem typically involved, our results indicate that these founder effects were

equally high irrespective of the age of the reservoirs. Several case studies on the propagule banks of *Daphnia* populations have demonstrated high local genetic stability over periods of 50–150 yr (Decaestecker et al. 2007; Mergeay et al. 2007). Recently, Ventura et al. (2014) even provided empirical evidence for founder effects lasting thousands of years. All this evidence indicates that zooplankton populations primarily have founder-controlled populations (Okamura and Freeland 2002), similar to founder-controlled communities (Sale 1977). In such populations, dispersal contributes little to gene flow and is mostly prevalent during the initial phase of colonization of empty or newly created habitats. While dormant propagules are the main unit of dispersal in most zooplankton, their most pervasive impact on landscape genetic structure may be their role in the short-term and long-term local persistence of populations as well as in fostering colonization of empty habitats rather than that they contribute to continuous gene flow among populations. Even seemingly extinct populations may still be recolonized by local dormant egg banks once the habitat becomes suitable again after decades (Mergeay et al. 2007). This has profound consequences for our view on metapopulation biology of zooplankton and other micro-organisms, as these species often share the lack of landscape genetic structure reflecting strong isolation-by-distance (Okamura and Freeland 2002). More specifically, we should not equal high potential for dispersal into high rates of gene flow (De Meester et al. 2016). In very small water bodies, however, negative effects of genetic drift and inbreeding can be pronounced, and the positive influence on fitness of immigrant alleles or genotypes from immigrants may then promote immigration and gene flow (Ebert et al. 2002). One might therefore expect a shift from a gene flow dominated system in extremely small populations (Ebert et al. 2002) to metapopulations that are more strongly dominated by local processes combined with extinction-recolonization dynamics in somewhat larger systems such as the reservoirs studied here, shallow lakes and the sometimes much smaller (approximately 100 m^2) farmland ponds (Vanoverbeke and De Meester 1997; De Meester et al. 2002; Louette et al. 2007).

Data availability statement

Microsatellite genotype data are stored in Dryad[®] under accession doi:10.5061/dryad.vh23q.

References

- Agostini, C., and others. 2010. Permanent genetic resources added to Molecular Ecology Resources Database 1 October 2010–30 November 2010. *Mol. Ecol. Resour.* **11**: 418–421. doi:10.1111/j.1755-0998.2010.02970.x
- Aguilera, X., J. Mergeay, A. Wollebrants, S. Declerck, and L. De Meester. 2007. Asexuality and polyploidy in *Daphnia* from the tropical Andes. *Limnol. Oceanogr.* **52**: 2079–2088. doi:10.4319/lo.2007.52.5.2079

- Asmelash, T., and others. 2007. Ecological atlas of reservoirs in Tigray, Northern Ethiopia. Tigray livelihood papers no. 4. VLIR-Mekelle Univ. IUC Programme and Zala-Daget Project.
- Bilton, D. T., J. R. Freeland, and B. Okamura. 2001. Dispersal in freshwater invertebrates. *Annu. Rev. Ecol. Syst.* **32**: 159–181. doi:10.1146/annurev.ecolsys.32.081501.114016
- Blanchet, F. G., P. Legendre, and D. Borcard. 2008. Forward selection of explanatory variables. *Ecology* **89**: 2623–2632. doi:10.1890/07-0986.1
- Boileau, M. G., P. D. N. Hebert, and S. S. Schwartz. 1992. Nonequilibrium gene-frequency divergence - persistent founder effects in natural populations. *J. Evol. Biol.* **5**: 25–39. doi:10.1046/j.1420-9101.1992.5010025.x
- Broquet, T., and E. J. Petit. 2009. Molecular estimation of dispersal for ecology and population genetics. *Annu. Rev. Ecol. Syst.* **40**: 193–216. doi:10.1146/annurev.ecolsys.110308.120324
- Cáceres, C. E., and D. A. Soluk. 2002. Blowing in the wind: A field test of overland dispersal and colonization by aquatic invertebrates. *Oecologia* **131**: 402–408. doi:10.1007/s00442-002-0897-5
- Cornuet, J. M., and G. Luikart. 1996. Description and power analysis of two tests for detecting recent population bottlenecks from allele frequency data. *Genetics* **144**: 2001–2014. PMC1207747.
- De Bie, T., and others. 2012. Body size and dispersal mode as key traits determining metacommunity structure of aquatic organisms. *Ecol. Lett.* **15**: 740–747. doi:10.1111/j.1461-0248.2012.01794.x
- De Meester, L., and J. Vanoverbeke. 1999. An uncoupling of male and sexual egg production leads to reduced inbreeding in the cyclical parthenogen *Daphnia*. *Proc. R. Soc. Lond. B Biol. Sci.* **266**: 2471–2477. doi:10.1098/rspb.1999.0948
- De Meester, L., A. Gómez, B. Okamura, and K. Schwenk. 2002. The Monopolization Hypothesis and the dispersal-gene flow paradox in aquatic organisms. *Acta Oecol.* **23**: 121–135. doi:10.1016/S1146-609X(02)01145-1
- De Meester, L., J. Vanoverbeke, K. De Gelas, R. Ortells, and P. Spaak. 2006. Genetic structure of cyclic parthenogenetic zooplankton populations – a conceptual framework. *Arch. Hydrobiol.* **167**: 217–244. doi:10.1127/0003-9136/2006/0167-0217
- De Meester, L., J. Vanoverbeke, L. J. Kilsdonk, and M. C. Urban. 2016. Evolving perspectives on monopolization and priority effects. *Trends Ecol. Evol.* **31**: 136–146. doi:10.1016/j.tree.2015.12.009
- Decaestecker, E., S. Gaba, J. A. M. Raeymaekers, R. Stoks, L. Van Kerckhoven, D. Ebert, and L. De Meester. 2007. Host-parasite ‘Red Queen’ dynamics archived in pond sediment. *Nature* **450**: 870–874. doi:10.1038/nature06291
- Dejenie, T., and others. 2008. Limnological and ecological characteristics of tropical highland reservoirs in Tigray, Northern Ethiopia. *Hydrobiologia* **610**: 193–209. doi:10.1007/s10750-008-9435-8
- Dray, S., P. Legendre, and P. R. Peres-Neto. 2006. Spatial modelling: A comprehensive framework for principal coordinate analysis of neighbour matrices (PCNM). *Ecol. Modell.* **196**: 483–493. doi:10.1016/j.ecolmodel.2006.02.015
- Ebert, D., C. Haag, M. Kirkpatrick, M. Riek, J. W. Hottinger, and V. I. Pajunen. 2002. A selective advantage to immigrant genes in a *Daphnia* metapopulation. *Science* **295**: 485–488. doi:10.1126/science.1067485
- Figuerola, J., and A. J. Green. 2002. Dispersal of aquatic organisms by waterbirds: A review of past research and priorities for future studies. *Freshw. Biol.* **47**: 483–494. doi:10.1046/j.1365-2427.2002.00829.x
- Foll, M., and O. Gaggiotti. 2006. Identifying the environmental factors that determine the genetic structure of populations. *Genetics* **174**: 875–891. doi:10.1534/genetics.106.059451
- Gu, Y., L. Xu, Q. Lin, H. Dumont, and B. Han. 2013. A new subspecies of *Daphnia*: *Daphnia similoides sinensis*. *Ecol. Sci.* **32**: 308–312. doi:10.1093/plankt/fbw031
- Haag, C. R., M. Riek, J. W. Hottinger, V. I. Pajunen, and D. Ebert. 2006. Founder events as determinants of within-island and among-island genetic structure of *Daphnia* metapopulations. *Heredity* **96**: 150–158. doi:10.1038/sj.hdy.6800774
- Haileselasie, T. H., J. Mergeay, L. J. Weider, R. Sommaruga, T. A. Davidson, M. Meerhoff, H. Arndt, K. Jürgens, E. Jeppesen, and L. De Meester. 2016. Environment not dispersal limitation drives clonal composition of Arctic *Daphnia* in a recently deglaciated area. *Mol. Ecol.* **25**: 5830–5842. doi:10.1111/mec.13843
- Hamilton, M. 2011. Population genetics. John Wiley & Sons.
- Hanski, I. 1998. Metapopulation dynamics. *Nature* **396**: 41–49. doi:10.1038/23876
- Haregeweyn, N., J. Poesen, J. Nyssen, J. De Wit, M. Haile, G. Govers, and S. Deckers. 2006. Reservoirs in Tigray (Northern Ethiopia): Characteristics and sediment deposition problems. *Land Degrad. Dev.* **17**: 211–230. doi:10.1002/ldr.698
- Hebert, P. D. N. 1987. Genotypic characteristics of cyclic parthenogens and their obligately asexual derivatives, p. 175–195. *In* S. C. Stearns [ed.], *The evolution of sex and its consequences*. Birkhäuser Basel.
- Jenkins, D. G., and A. L. Buikema. 1998. Do similar communities develop in similar sites? A test with zooplankton structure and function. *Ecol. Monogr.* **68**: 421–443. doi:10.1890/0012-9615(1998)068[0421:DSCDIS]2.0.CO;2
- Jost, L. 2007. Partitioning diversity into independent alpha and beta components. *Ecology* **88**: 2427–2439. doi:10.1890/06-1736.1
- Keenan, K., P. McGinnity, T. F. Cross, W. W. Crozier, and P. A. Prodöhl. 2013. diveRsity: An R package for the estimation and exploration of population genetics parameters and their associated errors. *Methods Ecol. Evol.* **4**: 782–788. doi:10.1111/2041-210X.12067
- Kumar, S., M. Nei, J. Dudley, and K. Tamura. 2008. MEGA: A biologist-centric software for evolutionary analysis of

- DNA and protein sequences. *Brief. Bioinform.* **9**: 299–306. doi:10.1093/bib/bbn017
- Legendre, P., and E. D. Gallagher. 2001. Ecologically meaningful transformations for ordination of species data. *Oecologia* **129**: 271–280. doi:10.1007/s004420100716
- Legendre, P., and M.-J. Fortin. 2010. Comparison of the Mantel test and alternative approaches for detecting complex multivariate relationships in the spatial analysis of genetic data. *Mol. Ecol. Resour.* **10**: 831–844. doi:10.1111/j.1755-0998.2010.02866.x
- Leibold, M. A., and others. 2004. The metacommunity concept: A framework for multi-scale community ecology. *Ecol. Lett.* **7**: 601–613. doi:10.1111/j.1461-0248.2004.00608.x
- Levins, R. 1969. Some demographic and genetic consequences of environmental heterogeneity for biological control. *Bull. Entomol. Soc. Am.* **15**: 237–240. doi:10.1093/besa/15.3.237
- Louette, G., and L. De Meester. 2005. High dispersal capacity of cladoceran zooplankton in newly created communities. *Ecology* **86**: 353–359. doi:10.1890/04-0403
- Louette, G., and L. De Meester. 2007. Predation and priority effects in experimental zooplankton communities. *Oikos* **116**: 419–426. doi:10.1111/j.2006.0030-1299.15381.x
- Louette, G., J. Vanoverbeke, R. Ortells, and L. De Meester. 2007. The founding mothers: The genetic structure of newly established *Daphnia* populations. *Oikos* **116**: 728–741. doi:10.1111/j.0030-1299.2007.15664.x
- Mergeay, J., J. Vanoverbeke, D. Verschuren, and L. De Meester. 2007. Extinction, recolonisation and dispersal through time in a planktonic crustacean. *Ecology* **88**: 3032–3043. doi:10.1890/06-1538.1
- Montero-Pau, J., A. Gómez, and J. Muñoz. 2008. Application of an inexpensive and high-throughput genomic DNA extraction method for the molecular ecology of zooplanktonic diapausing eggs. *Limnol. Oceanogr.: Methods* **6**: 218–222. doi:10.4319/lom.2008.6.218
- Nei, M., and R. K. Chesser. 1983. Estimation of fixation indices and gene diversities. *Ann. Hum. Genet.* **47**: 253–259. doi:10.1111/j.1469-1809.1983.tb00993.x
- Okamura, B., and J. R. Freeland. 2002. Gene flow and the evolutionary ecology of passively dispersing aquatic invertebrates, p. 194–216. *In* J. M. Bullock, R. E. Kenward, and R. S. Hails [eds.], *Dispersal ecology: 42nd Symposium of the British Ecological Society*. Cambridge University press, Cambridge.
- Orsini, L., J. Vanoverbeke, I. Swillen, J. Mergeay, and L. De Meester. 2013. Drivers of population genetic differentiation in the wild: Isolation-by-dispersal limitation, isolation-by-adaptation and isolation-by-colonization. *Mol. Ecol.* **22**: 5983–5999. doi:10.1111/mec.12561
- Pantel, J. H., T. E. Juenger, and M. A. Leibold. 2011. Environmental gradients structure *Daphnia pulex* × *pulicaria* clonal distribution. *J. Evol. Biol.* **24**: 723–732. doi:10.1111/j.1420-9101.2010.02196.x
- Popova, E. V., and others. 2016. Revision of the Old World *Daphnia* (Ctenodaphnia) *similis* group Cladocera: Daphniidae. *Zootaxa* **4161**: 1–40. doi:10.11646/zootaxa.4161.1.1
- R Core Team 2014. R: A language and environment for statistical computing (version 2.14). R Foundation for Statistical Computing, Vienna, Austria. Available from <http://www.R-project.org/>
- Rozas, J., J. C. Sanchez-DelBarrio, X. Messeguer, and R. Rozas. 2003. DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics* **19**: 2496–2497. doi:10.1093/bioinformatics/btg359
- Sale, P. F. 1977. Maintenance of high diversity in coral reef fish communities. *Am. Nat.* **111**: 337–359. doi:10.1086/283164
- Vanoverbeke, J., and L. De Meester. 1997. Among-population genetic differentiation in the cyclical parthenogen *Daphnia magna* (Crustacea, Anomopoda) and its relation to geographic distance and clonal diversity. *Hydrobiologia* **360**: 135–142. doi:10.1023/A:1003160903708
- Veech, J. A., and T. O. Crist. 2009. PARTITION: Software for hierarchical additive partitioning of species diversity, v. 3.0. Miami Univ.
- Ventura, M., A. Petrusek, A. Miró, E. Hamrová, D. Buñay, L. De Meester, and J. Mergeay. 2014. Local and regional founder effects in lake zooplankton persist after thousands of years despite high dispersal potential. *Mol. Ecol.* **23**: 1014–1027. doi:10.1111/mec.12656
- Wade, M. J., and D. E. McCauley. 1988. Extinction and recolonization: Their effects on the genetic differentiation of local populations. *Evolution* **42**: 995–1005. doi:10.2307/2408915
- Weir, B. S., and C. C. Cockerham. 1984. Estimating F-statistics for the analysis of population structure. *Evolution* **38**: 1358–1370. doi:10.1111/j.1469-1809.1949.tb02451.x
- Wright, S. 1949. The genetical structure of populations. *Ann. Eugenics* **15**: 323–354. doi:10.1111/j.14691809.1949.tb02451.x

Acknowledgments

The authors thank T. Dejenie, S. Risch, S. Declerck and others for help with sampling, and B. Hellemans, R. De Schutter, and S. Geldof for help with lab work. Two anonymous reviewers provided constructive and helpful comments on an earlier version. J. Mergeay was supported by a postdoctoral fellowship of the Research Foundation Flanders (FWO). This study was financially supported by a grant from the KULeuven Excellence Center (PF/2010/07). Samples were obtained thanks to financial support from the VLIR inter institutional cooperation (IUC) between Mekelle University (MU) and KU Leuven.

Conflict of Interest

None declared.

Submitted 24 August 2016

Revised 26 February 2017; 31 July 2017

Accepted 15 August 2017

Associate editor: Takehito Yoshida