

# Global change increases zoonotic risk, COVID-19 changes risk perceptions: a plea for urban nature connectedness

Maarten P.M. Vanhove , Séverine Thys , Ellen Decaestecker , Nicolas Antoine-Moussiaux , Jeroen De Man , Jean Hugé , Hans Keune , Ann Sterckx & Luc Janssens de Bisthoven

To cite this article: Maarten P.M. Vanhove , Séverine Thys , Ellen Decaestecker , Nicolas Antoine-Moussiaux , Jeroen De Man , Jean Hugé , Hans Keune , Ann Sterckx & Luc Janssens de Bisthoven (2020): Global change increases zoonotic risk, COVID-19 changes risk perceptions: a plea for urban nature connectedness, *Cities & Health*, DOI: [10.1080/23748834.2020.1805282](https://doi.org/10.1080/23748834.2020.1805282)

To link to this article: <https://doi.org/10.1080/23748834.2020.1805282>



Published online: 02 Sep 2020.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

## Global change increases zoonotic risk, COVID-19 changes risk perceptions: a plea for urban nature connectedness

Maarten P.M. Vanhove<sup>a,b</sup>, Séverine Thys<sup>c</sup>, Ellen Decaestecker<sup>d</sup>, Nicolas Antoine-Moussiaux<sup>e</sup>, Jeroen De Man<sup>f</sup>, Jean Hugé<sup>g,h,i</sup>, Hans Keune<sup>j,k</sup>, Ann Sterckx<sup>k</sup> and Luc Janssens de Bisthoven<sup>l</sup>

<sup>a</sup>Centre for Environmental Sciences, Research Group Zoology: Biodiversity and Toxicology, Hasselt University, Diepenbeek, Belgium; <sup>b</sup>Department of Botany and Zoology, Faculty of Science, Masaryk University, Brno, Czech Republic; <sup>c</sup>Department of Vaccinology, Faculty of Medicine and Health Sciences, University of Antwerp, Antwerp, Belgium; <sup>d</sup>Aquatic Biology, Interdisciplinary Research Facility Life Sciences, KU Leuven, Kortrijk, Belgium; <sup>e</sup>Fundamental and Applied Research for Animals and Health (FARAH), Faculty of Veterinary Medicine, University of Liège, Liège, Belgium; <sup>f</sup>Centre for General Practice, Department of Primary and Interdisciplinary Care, Faculty of Medical and Health Sciences, University of Antwerp, Antwerp, Belgium; <sup>g</sup>Department of Environmental Sciences, Open University of the Netherlands, Heerlen, The Netherlands; <sup>h</sup>Biology Department, Vrije Universiteit Brussel, Brussels, Belgium; <sup>i</sup>Centre for Environmental Sciences, Research Group Environmental Biology, Hasselt University, Diepenbeek, Belgium; <sup>j</sup>Belgian Biodiversity Platform – Team Nature and Society, Research Institute Nature and Forest (INBO), Brussels, Belgium; <sup>k</sup>Chair Care and the Natural Living Environment, Department of Primary and Interdisciplinary Care Antwerp (ELIZA), Faculty of Medical and Health Sciences, University of Antwerp, Antwerp, Belgium; <sup>l</sup>Capacities for Biodiversity and Sustainable Development (CEBioS), Royal Belgian Institute for Natural Sciences, Brussels, Belgium

### ABSTRACT

Ebola and COVID-19 are textbook emerging diseases influenced by humans. Ebola is often considered a result of exotic nature threatening health. Conversely, COVID-19, emerged in an urban environment, entails risks worldwide. Geographical, virological and demographic differences influence risk perceptions and responses to both diseases. Because ecological understanding of urban human-animal relations improves disease risk assessment, we call for ethnographical exploration of this interface. 'Global Urban Confinement Measures' impact health by influencing disease perceptions, limiting nature access, and strengthening inequities. To prevent and mitigate zoonotic pandemics and their consequences, policy should promote nature connectedness, concert with stakeholders, and integrate nature-city-inhabitant interactions.

### ARTICLE HISTORY

Received 5 May 2020  
Accepted 13 July 2020

### KEYWORDS

Eco Health; One Health; urban environmental justice


### Different risk perceptions with different zoonoses: towards 'shared risk'?

While COVID-19 dominates global news, comparisons are made with another well-known zoonosis: 'Ebola Virus Disease (EVD)', especially its 2014 West African episode. Like EVD, COVID-19 originated from viral spill-over from animals, but it resulted in a totally different epidemiology and societal response. As starting point for reconsidering the role of nature in cities, we focus on the striking extent to which COVID-19 surpasses EVD in global economic impact and societal response. The central position of China's economy, with interlinkages worldwide, has been proposed as a driver of this phenomenon (Maffioli 2020). We argue that their different geographic origins and setting also mediate response in policy and urban society (Figure 1), and expect debate on underlying mechanisms to continue.

In this argument, risk framing for adequate policy response is crucial, both by authorities (Peeri *et al.* 2020), and by the public (Betsch 2020, Lohiniva *et al.* 2020). Despite between-country differences (e.g. less strict containment, more attention for testing and contact tracing in South Korea: Tanne *et al.* 2020) more and more governments deemed risk levels sufficiently high to

implement often unprecedented travel restrictions and population 'lockdown', initially focusing on cities (see Lau *et al.* 2020). Insofar as they are state-imposed and regard cities, we here refer to these public health actions as 'Global Urban Confinement Measures (GUCMs)'. They are considered to have contributed to mitigating the COVID-19 pandemic (e.g. Denis *et al.* 2020, Lau *et al.* 2020). Moreover, in data-poor conditions, as is often the case with little understood diseases or in the Global South, behavioural changes, combined with detection and surveillance, are crucial to reduce epidemic transmission in urban areas (Moss *et al.* 2016).

Although the practice and logic of confinement are ancient and partly similar actions were applied during EVD outbreaks (Peak *et al.* 2018), today's GUCMs render their effects tangible to everyone because they are per definition ubiquitous. Thus, health-related and other (economic, social) risks associated with COVID-19 and future emerging infectious diseases are currently hard to downplay, as they will directly influence control, preventive and reactive strategies in different countries. Next to contrasting media uproar, we also observe stark differences in risk perceptions underlying the responses to



**Figure 1.** Scheme of how perceptions of nature, and land use at the human-animal-environment interface (horizontal axis) and the spatial scale of mobility (vertical axis) may relate to the intensity and scope of the policy response (diagonal axis). This is exemplified for the new human-animal-environmental interactions that constituted the outbreak contexts of mainly rural Ebola Virus Disease (EVD) and mainly urban COVID-19. The three axes and four quadrants demonstrate the inter- and transdisciplinarity needed to characterize the complexity of pandemics in (urban) society, and the positions of elements within this scheme remains open for interpretation. We propose that reconnection with nature at various levels may be a response mitigating the effects of the pandemic and of the policy response to it.

EVD and COVID-19, respectively. Like multiple other recent zoonotic outbreaks, and certainly as it hardly directly affected the Global North, public opinion may consider EVD restricted to ‘exotic’, often rural places where (proximity to) nature seems antagonistic to human health (Antoine-Moussiaux *et al.* 2019). This stance focuses on animals as a health threat to humans: ‘us *versus* them’. It contrasts with a ‘shared risk’ perspective, in which animals are models and sentinels for understanding infectious disease and its environmental drivers, because humans and non-human animals share environmental and infection hazards (Rabinowitz *et al.* 2008). By contrast, COVID-19’s societal impact is unseen in recent history: it is not limited to the Global South, and originated in Wuhan, an urban centre of economic activity. Transport duration of bushmeat, remoteness and low human population density limited chances of spreading EVD overseas, as it mainly depends on direct contact with bodily fluids of patients (Kümpel *et al.* 2015). Conversely, Wuhan’s strong inter-continental transport links, and COVID-19’s

contagiousness (Peeri *et al.* 2020) prompted widespread adoption of GUCMs to avoid COVID-19 overrunning health systems (Figure 1). As such, GUCMs are both a crisis response, and an epiphenomenon confronting (predominantly) urban communities with a cosmopolitan vulnerability to zoonoses.

Even though proximity to nature is often low, the threat of zoonoses may create perceptions to further limit human-nature interactions in urban areas. We argue that this would be a mistake and render GUCMs unsustainable. On the contrary, COVID-19 should spur urban populations worldwide, often physically and mentally separated from nature, to also see the human-animal-environment nexus from a ‘shared risk’ perspective. Integrative approaches such as One Health, EcoHealth, and Planetary Health, jointly consider the health of humans, animals and the environment. These frameworks make the connections between the impacts on these three domains of health explicit and promote a holistic approach of the human-animal-ecosystem health nexus at the science-policy interface (Keune

*et al.* 2017). From a One Health perspective, we argue that the COVID-19 crisis gives the opportunity to consider these three research levels: (1) the study of ecological and evolutionary dynamics mediating disease transmission in urban environments, (2) the ethnography of individual interactions between humans and urban biodiversity, and (3) the community psychology of nature's role in public health and urban space.

### The human-animal-environment interface in urban evolutionary ecology of disease

Within the human-animal-environment interface, anthropogenic landscape and habitat changes, as well as unregulated animal exploitation, facilitate zoonotic viral transmission (Johnson *et al.* 2020, Pratesi 2020) (Figure 1). Cause-effect relationships underlying zoonoses are however not straightforward; for example, highest transmission risks were suggested to occur at intermediate levels of land use change (Faust *et al.* 2018). Also, human-induced environmental and physiological stress in wildlife alter its pathogen dynamics (Plowright *et al.* 2015, Seltmann *et al.* 2017, Rohr *et al.* 2019 and references therein). Urbanisation may be another factor altering wildlife pathogen dynamics (Fountain-Jones *et al.* 2017) and increasing viral spill-over (Smiley Evans *et al.* 2020), hence the particular relevance to cities. Potential examples of new human-animal-environment interfaces (Figure 1) include increased human-animal contact through informal or low-biosecurity livestock keeping or wet markets, and unintended escape from laboratories close to city centres (examples in Houwenhuysen *et al.* 2018). Also, urban animal populations differ from rural conspecifics in microbial (Flandroy *et al.* 2018) and parasite communities (Rouffaer *et al.* 2017), which may affect the hosts' immunity. Unsurprisingly, the relatively recent discipline of urban ecology, intersecting with evolutionary mechanisms in the emerging field of 'urban evolutionary ecology', is relevant to urban planning and human public health (Rivkin *et al.* 2019). Given the complex interplay between these host-related and environmental factors, we cannot agree more with Trinh *et al.* (2018) as they state that 'One Health approaches will benefit from ecological and evolutionary studies that distill natural variation in host-microbe interaction into a more tractable number of general rules'. Because of these host-related and environmental factors, and given an adequate science-policy interface, translating urban ecology into urban design and planning could be one of the many possibilities for research supporting public health decision-making (see Sallis *et al.* 2016). There are strong indications that connectedness (both mental and physical) with, and awareness about nature are related to nature's health benefits (Frumkin *et al.* 2017). Therefore, we wonder whether better comprehension of the human-animal-environment interface is also

beneficial to human and environmental health, which would be complementary to the purely practical advantage of better-informed public health policy response. Still, all early warning systems failed to prevent or contain the COVID-19 epidemic. The reasons why still need to be analysed in depth, but scientific awareness and understanding of the medical and veterinary relevance of these biological interactions will clearly not suffice in avoiding and tackling urban zoonotic outbreaks and their effects (see the phenomenon of 'late lessons from early warnings' in the field of risk assessment: EEA 2013). We have no indications that people's behavioural compliance to GUCMs implies awareness of, let alone changed attitudes towards, urban ecology at the human-animal-environment nexus. We suggest this crisis should therefore serve as an occasion to rethink the role of nature in the city and vice-versa at two other levels, too: the shared interests of urban dwellers and non-human animals, and the community psychology of urban nature and health.

### Human animal health in medical anthropology

The role of non-human animals in urban health is multifaceted. Animals can be hosts to zoonotic diseases, with COVID-19 as compelling example of the impact of bushmeat trade and consumption in cities. Animals can also be disease vectors, e.g. for dengue or malaria, as mosquitoes find good breeding grounds under urban conditions. In this context, Kelly and Lezaun (2014) explain that public health interventions often aim at 'interspecies distancing' including extinction. (At present it is hard not to draw a parallel with ongoing COVID-19-related GUCMs.) Urban planning therefore influences connections and cohabitation between humans and mosquitoes and other animals in their 'shared built environment' (see Kelly and Lezaun 2014). As mentioned above, insight into the ecology of disease has management applications. It allows, e.g. vector control to not only rely on insecticides but also human behavioural changes, vector modification, improved housing, or removal of trash that offers mosquito breeding grounds (Arunachalam *et al.* 2012, Finda *et al.* 2020, Forsyth *et al.* 2020). Given adequate awareness and scientific understanding, we therefore propose to reconsider cohabitation with pathogens instead of eradicating them within fragile 'ultra-hygienic societies'. Recent developments in this context include the Microbial Theory of Health, encouraging 'targeted hygiene' focusing only on high-risk sites, surfaces and practices for pathogen transmission (Scott *et al.* 2020); the medical applications of the Hygiene Hypothesis on the positive immunoregulatory role of certain pathogens (Versini *et al.* 2015); and the Microbiome Rewilding Hypothesis. The latter hinges on the link between biodiversity loss and altered microbial exposure leading to immune dysregulation among urban populations. The hypothesis

proposes that restoring urban biodiversity may therefore rewild environmental microbiomes in urban nature. They could then benefit human health through immune protection and hence disease prevention (Mills *et al.* 2017). We suggest to investigate how urban space can be redesigned for restorative urban gardens that benefit human well-being (*sensu* Thwaites *et al.* 2005) but also create space for cohabitation with animals and environmental rewilding. Since anthropogenic pressures obviously impact the risk originating from human-animal contact, the above-mentioned ‘shared risk’ perspective becomes all the more relevant. Indeed, public health policy should look beyond managing animals to benefit human welfare and disease control. The complex dynamics of human-animal encounters require a ‘multispecies ethnography’ (Brown and Nading 2019). A practical example could be changing the existing zoonotic cycles diagram used in life sciences and public health to visualize infections with both animal and human hosts. These diagrams far from simply summarize the epidemiological reality of human-animal transmission. They are unique ways of sketching forms, patterns, and relations, visualising different aspects of interspecies existence. In their graphical choices, they convey a peculiar vision of health and nature, often translating a myth of mastery of nature by humans (Lynteris 2017). This means they are most often centred around the survival of the human species (over the non-human species), representing wildlife as a threat, and showing traditional rural customs as a cause for transmission (Thys 2019a). This visual framing, entering the realm of the ‘us *versus* them’ thinking, should be questioned and modified. This calls for research to better understand how visual representations might support ‘shared risk’ framing. We therefore ask more attention for an emerging medical-anthropological field: ‘human animal health’ (Brown and Nading 2019).

Moving beyond animals and towards urban green spaces at societal level, we witnessed that COVID-19 and its lockdown scenarios highlighted the role of urban nature in human equity and well-being. Importantly for urban planners and other policymakers, urban nature, apart from its health benefits, is also linked to beneficial environmental effects relevant at abovementioned levels, such as species conservation (Hartig *et al.* 2014, Helm 2020).

### **An environmental health and ecopsychological lens on COVID-19 impacts on urban individuals and societies**

We think the many beneficial effects of nature exposure and reconnection with nature are strongly underestimated (Figure 1). They include children’s academic achievement and healthy development (Kuo *et al.* 2019), cognitive functioning, emotional well-being, and other dimensions of mental health (Bratman *et al.* 2019).

Moreover, nature connectedness, in the sense of a persistent awareness of the interrelatedness between one’s self and nature, correlates positively with sustainable behaviour (Zylstra *et al.* 2014, Whitburn *et al.* 2019). While GUCMs force human beings to reshape their relation amongst themselves and with urban space and nature, such measures proved poorly inclusive towards the urban population’s entire socioecological diversity (*sensu* Greenaway and Turetsky 2020, i.e. with respect to the distribution of social groups under given settings of the physical, political or social environment). GUCMs aggravate unequal access to urban nature which undermines their acceptability. Disadvantaged slum populations with poor housing, limited access to water and mobility and high population density typically have poor access to nature in their close proximity, potentially further compounded by GUCMs. Also, avoiding human gathering in public spaces may result in a proportionally bigger loss among slum dwellers since they typically spend more time outside and are practically unable to abide to confinement rules. This may change their lives from difficult to unbearable. COVID-19 also underscores another perspective of human inequity regarding nature and urban space. Disadvantaged populations are disproportionately affected by various communicable and non-communicable conditions (e.g. diabetes, hypertension, likely also tuberculosis) making them more vulnerable to COVID-19-induced morbidity and mortality (e.g. Boffa *et al.* 2020, Denis *et al.* 2020). Such conditions often thrive in disadvantaged and more densely populated urban areas because of a variety of reasons, including closer social and physical contact among humans, lack of space/nature to perform physical activity or to grow fresh and healthy food, increased promotion of and access to processed food, etc. (De Man *et al.* 2019). Inequity from pre-existing comorbidities is likely compounded even more through positive feedback loops. Indeed, ‘collateral damage’ results from higher risks of nosocomial (hospital-acquired) COVID-19 transmission among people attending health services, fear-induced decrease in health-seeking behaviour and decreased health service availability due to COVID-19. Thus, GUCMs defy urban environmental justice in various ways. When aiming at urban environmental justice, the planning of urban green spaces needs to be considered together with aspects of citizens’ health and their social networks, within the framework of the city as an integrated socio-ecological system (Enssle and Kabisch 2020). Indeed, approaching human, animal and environmental health in an integrative way may contribute to environmental justice (Rüegg *et al.* 2018). Without this integration, a disconnection of nature follows, and we propose that GUCMs in that case aggravate an already major current public health challenge (see Anguelovski 2016, Corburn 2017).

We suggest this indicates insufficient dialogue about the potential effects of GUCMs amongst

citizens, civil society and decision-makers (including parliaments), and a lack of translation by decision-makers of expert recommendations. Indeed, expert-derived information requires stakeholder involvement and shared problem framing to be rendered into so-called socially robust knowledge (Cornell *et al.* 2013). Knowledge integration and co-production are indeed necessary for any One Health approach to be successfully implemented into policy (Rüegg *et al.* 2018). Without societal debate, we expect societal response to health challenges like COVID-19 to be unsustainable, as GUCMs exacerbate existing inequities. Compliance to GUCMs is however a combined product of agency of individuals and communities, urging us for a community psychology perspective. GUCMs remove citizens' sense of individual control. Resulting isolation can cause or worsen negative feelings (e.g. loneliness, anxiety, depression symptoms). This makes people ill at ease in cities in times of COVID-19. In contrast, during EVD outbreaks different reaction dynamics occurred, as movement from rural to urban areas was observed (Fallah *et al.* 2018). We question how urban populations feel about the trade-off between the benefits of GUCMs of lowering infection risk, and their negative consequences. Also here, considering a less anthropocentric, non-'us versus them' stance is worthwhile: exposure to urban nature has proven mental health benefits (e.g. through psychological restoration: Hartig *et al.* 2014). It may contribute to reduction of health inequalities (WHO 2016) e.g. through healthier behaviour (e.g. physical activity) that would limit above-mentioned co-morbidities, and may contribute to behavioural changes towards environmental sustainability (Nisbet and Zelenski 2011, Ives *et al.* 2018, Whitburn *et al.* 2019). We hypothesize a positive feedback loop whereby more access to urban nature forcibly close by (due to GUCMs) mitigates GUCMs' negative effects, but also makes city dwellers' attitudes less prone to causing environmental impact, e.g. habitat degradation, contributing to future pandemics. For example, we expect well-distributed, attractive urban green to promote outdoor leisure, contributing to health and pro-environmental behaviour. We therefore propose that ensuring sufficient access to (urban) nature and exploring its ecopsychological aspects (see Fisher 2013, Roszak *et al.* 1995, Vakoch and Castrillón 2014) will help rendering potential future GUCMs inclusive, educational, sustainable and transformational. An avenue for implementation may be urban agriculture, which provides diverse ecosystem services to urban dwellers, society and the environment (Cartiaux *et al.* 2018).

We indeed cannot predict the potential sustainability benefits of GUCMs. Relevant to the urban

human-animal-environment nexus are increased wildlife sightings (Bates *et al.* 2020) and potentially higher valuation of urban nature (Helm 2020). Across continents, local improvements in urban air quality were observed, exemplified by reports from Milan, Italy (Collivignarelli *et al.* 2020), Rio de Janeiro, Brazil (Dantas *et al.* 2020) and Salé, Morocco (Otmani *et al.* 2020). However, these studies also mention stark contrasts between pollutants, and between the city and its surroundings because of influences from adjacent areas with different land use. Also, already in China, these positive environmental impacts are expected to be temporary (Wang and Su 2020). Worldwide, COVID-19-related decreases in, e.g. conservation management, conservation enforcement, scientific attention or ecotourism revenue may have adverse environmental impacts (Bates *et al.* 2020, Helm 2020).

### Perspectives for policy extension across North and South: not a one-stop-shop

There are indications that human psychology is adapted to local disease ecology situations: collectivism may work against infections under high pathogen prevalence (Thys 2019b and references therein), encouraging more international comparative approaches to infectious disease and health policy. While we focus here on environmental and psychosocial aspects of health, it merits investigation whether a 'shared risk' perspective can also foster cross-cultural or cross-national comparisons inspiring policy response to COVID-19 and other zoonoses. What COVID-19 will mean for cities in the Global South is as yet hard to fathom, with their sharply visible inequities experienced by urban slum populations, and with transmission noticeably from well-to-do travellers to poorer communities (e.g. Manderson and Levine 2020). As the Global South accounts for the fastest urbanisation worldwide, and with that, most of global demographic growth, its cities hold great interest and potential for sustainability; regrettably, they are underrepresented in research (Nagendra *et al.* 2018). We cannot stress enough that GUCMs are clearly no one-stop-shop implementable in an identical way in the Global North and South alike (see also Manderson and Levine 2020). While lockdown measures have been implemented throughout countries that strongly differ social-politically and geographically, also within the North strategies for surveillance and reduction of transmission vary widely between states, among others influenced by resource availability, culture, governance and legislation (Cohen and Kupferschmidt 2020). New Zealand scientists advised their government that looking at responses in a set of Asian countries may be informative (Wilson *et al.* 2020). Although COVID-19 may have simply arrived late in Africa and there is no certainty whether the African epidemics will remain

small (Martinez-Alvarez *et al.* 2020), COVID-19 seemed to progress slower in Africa, and important lessons have been learnt from EVD (Nuwagira and Muzoora 2020). We therefore hope that best practices from this pandemic may inspire urban policies cross-continently. We also hope that instead of seeing nature as a threat, more nature connectedness and less anthropocentric interactions with nature may render future such policies more sustainable (Figure 1). For example, given the zoonotic importance of human-animal-environment interactions and the potential of animal disease models, better understanding of hosts, vectors and parasites necessitates additional research. The call for multispecies ethnography in human animal health reflects the need to grant urban animals and nature space to avoid transmission in new human-animal-environment interfaces. These efforts in urban ecology and medical anthropology would in our view contribute to pandemic prevention. Lastly, the mental and physical nature health benefits extend not only to avoiding pandemics but also to reacting to them and their consequences. Therefore, to mitigate harmful effects of future pandemics and the ensuing GUCMs, we urge urban planners and other policymakers to ensure equitable access to urban green.

## Acknowledgments

Capacity development and bridge building within the Belgian One Health Network ([www.biodiversity.be/onehealth](http://www.biodiversity.be/onehealth)) facilitated the preparation of this commentary. Lucie Ongena (Royal Belgian Institute of Natural Sciences) is cordially thanked for providing Figure 1, Nikol Kmentová (Masaryk University/Hasselt University) and Armando J. Cruz-Laufer (Hasselt University) for suggestions on Figure 1, and Leila Demarest (Leiden University) for thorough feedback on the manuscript. The comments of anonymous reviewers are gratefully acknowledged.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Notes on contributors

**Maarten P. M. Vanhove**, biologist and social-cultural anthropologist, holds a PhD in Sciences (KU Leuven). He was researcher at the Royal Museum for Central Africa, the Hellenic Centre for Marine Research, and Masaryk University. Subsequently, he worked on capacity development as a biodiversity policy scientist at the Royal Belgian Institute of Natural Sciences, before becoming curator of worms at the Finnish Museum of Natural History within the University of Helsinki. Currently, he is assistant professor in Zoology/Aquatic Biodiversity at Hasselt University. He studies biodiversity policy, and parasite genetics and morphology in evolutionary and conservation-relevant contexts, focusing on the Mediterranean and Africa.

**Séverine Thys**, medical anthropologist, master in Public Health (ULB-ULg-UCL) and PhD in Veterinary Sciences (UGhent), was researcher at the Institute of Tropical Medicine (Antwerp), on perceptions of zoonotic diseases in African and Asian rural communities. She coordinated a strategic network to foster intersectoral collaboration and promote the 'One Health' concept (Belgian Development Cooperation) and was co-investigator in three FP7 EU projects (ICONZ, ADVANZ, NEXTGEN). She was deployed in Guinea for GOARN (WHO) during the EVD epidemic in West Africa and works now at the Centre for Evaluation of Vaccinations and the Global Health Institute at UA for the EBOVAC3 project.

**Ellen Decaestecker**, biologist, was postdoctoral research fellow at the Centre of Immunity, Infection and Evolution (Edinburgh University). She is professor (KU Leuven, Biology Department) and teaches Ecology, Evolution, Comparative, Human and Zoological Biology and Animal Diversity. She is president of the Royal Belgian Zoological Society, member of the KU Leuven Research Council, FWO BIO3 panel and was scientific member of the Belgian Governmental Committee Health and Environment. She engages in societally relevant science, urgently needed to solve problems associated with global change and sustainability. She is specialized in host-microbiome interactions with a focus on host-parasite coevolution.

**Nicolas Antoine-Moussiaux**, doctor in veterinary medicine, holds a PhD in veterinary sciences (University of Liège). After a master in Rural sociology and economics (Catholic University of Louvain/Gembloux Agro-Bio tech), he engaged into an interdisciplinary project about surveillance of avian influenza in Vietnam with CIRAD (French institute of research for international agricultural development), while still pursuing a research agenda about animal genetic resources management in Africa. On this basis, as senior lecturer at University of Liège, he has developed a master program to operationalize One Health as integrated management of health risks, and a research thread along the same line.

**Jeroen De Man**, a medical doctor specialized in primary care (KU Leuven), holds a master's in global public health policy (UNC-Chapel Hill) and a postgraduate certificate in tropical medicine and international health (ITM, Antwerp). Jeroen was working as a family doctor in a disadvantaged area in Brussels. He lived and worked in rural India to strengthen the local health system. He was a researcher at the Institute of Tropical Medicine in Antwerp. Currently employed by the University of Antwerp, he lives in Cape Town where he studies health behavior among people with diabetes living in urban slums.

**Jean Hugé**, sustainability scientist, holds a Master in BioScience engineering and in Conflict and Development (Ghent University) and a PhD in Sciences (Vrije Universiteit Brussel and Université Libre de Bruxelles). He is currently Assistant Professor in Environmental Governance at the Open University of the Netherlands, and also teaches at Hasselt University and Vrije Universiteit Brussel. He studies sustainability assessment, conservation conflicts, biodiversity governance and participatory natural resources management, with a focus on coastal tropical ecosystems.

**Hans Keune**, political scientist (University of Amsterdam) with a PhD in Environmental Sciences (University of Antwerp), works on critical complexity, OneHealth/

EcoHealth. Works at the Belgian Biodiversity Platform, the Research Institute for Nature and Forest (INBO) and is professor at the Faculty of Medicine and Health Sciences of the University of Antwerp where he coordinates the Chair Care and the Natural Living Environment. Coordinated the organization of the European OneHealth/EcoHealth workshop (Brussels, Oct. 2016) and co-coordinates the Belgian One Health Network. Was Lead Author on the IPBES Regional Assessment Europe – Central Asia where he mainly contributed on health.

**Ann Sterckx**, ecopsychologist, holds a Master in Organizational Psychology (Ghent University). In her work she specialised in Ecopsychology and Ecotherapy. Currently she runs EarthWise Education, a Learning Centre for ecopsychology and nature connectedness. Ann is an active member of the global International Ecopsychology Society. Since 2019 she is PhD candidate at the Faculty of Medicine and Health Sciences of the University of Antwerp, in the framework of the Chair of Care and the Natural Living Environment. Her PhD research focuses on the relationship between nature connectedness, self-care and professional practice.

**Luc Janssens de Bisthoven**, biologist, holds a PhD in Sciences (KU Leuven) as aquatic ecotoxicologist. He was lecturer at Kenyatta University and researcher at Lund University, Rhodes University, University of Aveiro and Beijing University. He worked as independent consultant on water quality surveillance in Germany and as programme officer at the Flemish Council for InterUniversity Development Cooperation (VLIR-UOS) with focus on East Africa and Suriname. Now he is guest speaker at VUB and UHasselt and coordinator of CEBioS at the Royal Belgian Institute of Natural Sciences, on capacity building in developing countries on biodiversity conservation and policy, with focus on Benin, Burundi, DR Congo and Vietnam.

## ORCID

Maarten P.M. Vanhove  <http://orcid.org/0000-0003-3100-7566>

Séverine Thys  <http://orcid.org/0000-0001-7936-3194>

Ellen Decaestecker  <http://orcid.org/0000-0001-6328-5283>

Nicolas Antoine-Moussiaux  <http://orcid.org/0000-0003-1575-625X>

Jeroen De Man  <http://orcid.org/0000-0002-3196-8757>

Jean Hugé  <http://orcid.org/0000-0002-3695-547X>

Hans Keune  <http://orcid.org/0000-0002-1531-944X>

Luc Janssens de Bisthoven  <http://orcid.org/0000-0002-9547-1563>

## References

- Anguelovski, I., 2016. From toxic sites to parks as (green) LULUs? New challenges of inequity, privilege, gentrification, and exclusion for urban environmental justice. *Journal of planning literature*, 31 (1), 23–36. doi:10.1177/0885412215610491
- Antoine-Moussiaux, N., et al., 2019. The good, the bad and the ugly: framing debates on nature in a one health community. *Sustainability science*, 14 (6), 1729–1738. doi:10.1007/s11625-019-00674-z
- Arunachalam, N., et al., 2012. Community-based control of *Aedes aegypti* by adoption of eco-health methods in Chennai City, India. *Pathogens and global health*, 106 (8), 488–496. doi:10.1179/2047773212Y.0000000056
- Bates, A.E., et al., 2020. COVID-19 pandemic and associated lockdown as a “Global human confinement experiment” to investigate biodiversity conservation. *Biological conservation*, 108665. doi:10.1016/j.biocon.2020.108665
- Betsch, C., 2020. How behavioural science data helps mitigate the COVID-19 crisis. *Nature human behaviour*, 4, 438. doi:10.1038/s41562-020-0866-1
- Boffa, J., et al., 2020. COVID-19 and tuberculosis in South Africa: A dangerous combination. *SAMJ: South African medical journal*, 110 (5), 1–2. doi:10.7196/SAMJ.2020.v110i5.14747
- Bratman, G.N., et al., 2019. Nature and mental health: an ecosystem service perspective. *Science advances*, 5 (7), eaax0903. doi:10.1126/sciadv.aax0903
- Brown, H. and Nading, A.M., 2019. Introduction: human animal health in medical anthropology. *Medical anthropology quarterly*, 33 (1), 5–23. doi:10.1111/maq.12488
- Cartiaux, N., et al., 2018. Improving urban metabolism through agriculture: an approach to ecosystem services qualitative assessment in Rome. *Vertigo Hors-série*, 31. doi:10.4000/vertigo.21655
- Cohen, J. and Kupferschmidt, K., 2020. Countries test tactics in ‘war’ against COVID-19. *Science*, 367 (6484), 1287–1288. doi:10.1126/science.367.6484.1287
- Collivignarelli, M.C., et al., 2020. Lockdown for CoViD-2019 in Milan: what are the effects on air quality? *Science of the total environment*, 732, 139280. doi:10.1016/j.scitotenv.2020.139280
- Corburn, J., 2017. Urban place and health equity: critical issues and practices. *International journal of environmental research and public health*, 14 (2), 117. doi:10.3390/ijerph14020117
- Cornell, S., et al., 2013. Opening up knowledge systems for better responses to global environmental change. *Environmental science & policy*, 28, 60–70. doi:10.1016/j.envsci.2012.11.008
- Dantas, G., et al., 2020. The impact of COVID-19 partial lockdown on the air quality of the city of Rio de Janeiro, Brazil. *Science of the total environment*, 729, 139085. doi:10.1016/j.scitotenv.2020.139085
- De Man, J., et al., 2019. Diabetes self-management in three different income settings: cross-learning of barriers and opportunities. *Plos one*, 14 (3), e0213530. doi:10.1371/journal.pone.0213530
- Denis, M., et al., 2020. COVIPENDIUM: information available to support the development of medical countermeasures and interventions against COVID-19. *Transdisciplinary Insights Living Paper*. Leuven: KU Leuven. Available at: [https://rega.kuleuven.be/if/corona\\_covid-19](https://rega.kuleuven.be/if/corona_covid-19)
- Enssle, F. and Kabisch, N., 2020. Urban green spaces for the social interaction, health and well-being of older people – an integrated view of urban ecosystem services and socio-environmental justice. *Environmental science & policy*, 109, 36–44. doi:10.1016/j.envsci.2020.04.008
- European Environment Agency, 2013. *Late lessons from early warnings: science, precaution, innovation – Summary*. EEA Report 1/2013. Luxembourg: Publications Office of the European Union. Available from: <https://www.eea.europa.eu/publications/late-lessons-2>
- Fallah, M.P., Skrip, L.A., and Enders, J., 2018. Preventing rural to urban spread of Ebola: lessons from Liberia. *The lancet*, 392 (10144), 279–280. doi:10.1016/S0140-6736(18)31435-1
- Faust, C.L., et al., 2018. Pathogen spillover during land conversion. *Ecology letters*, 21 (4), 471–483. doi:10.1111/ele.12904

- Finda, M.F., *et al.*, 2020. Opinions of key stakeholders on alternative interventions for malaria control and elimination in Tanzania. *Malaria journal*, 19, 164. doi:10.1186/s12936-020-03239-z
- Fisher, A., 2013. *Radical ecopsychology: psychology in the service of life*. Albany: SUNY Press.
- Flandroy, L., *et al.*, 2018. The impact of human activities and lifestyles on the interlinked microbiota and health of humans and of ecosystems. *Science of the total environment*, 627, 1018–1038. doi:10.1016/j.scitotenv.2018.01.288
- Forsyth, J.E., *et al.*, 2020. Source reduction with a purpose: mosquito ecology and community perspectives offer insights for improving household mosquito management in coastal Kenya. *PLoS neglected tropical diseases*, 14 (5), e0008239. doi:10.1371/journal.pntd.0008239
- Fountain-Jones, N.M., *et al.*, 2017. Urban landscapes can change virus gene flow and evolution in a fragmentation-sensitive carnivore. *Molecular ecology*, 26 (22), 6487–6498. doi:10.1111/mec.14375
- Frumkin, H., *et al.*, 2017. Nature contact and human health: A research agenda. *Environmental health perspectives*, 125 (7), 075001. doi:10.1289/EHP1663
- Greenaway, V.P. and Turetsky, K.M., 2020. Socioecological diversity and inclusion: a framework for advancing diversity science. *Current opinion in psychology*, 32, 171–176. doi:10.1016/j.copsy.2019.09.008
- Hartig, T., *et al.*, 2014. Nature and Health. *Annual review of public health*, 35 (1), 207–228. doi:10.1146/annurev-publhealth-032013-182443
- Helm, D., 2020. The environmental impacts of the coronavirus. *Environmental & resource economics*, 76, 21–38. doi:10.1007/s10640-020-00426-z
- Houwenhuysse, S., *et al.*, 2018. Back to the future in a petri dish: origin and impact of resurrected microbes in natural populations. *Evolutionary applications*, 11 (1), 29–41. doi:10.1111/eva.12538
- Ives, C.D., *et al.*, 2018. Reconnecting with nature for sustainability. *Sustainability science*, 13, 1389–1397. doi:10.1007/s11625-018-0542-9
- Johnson, C.K., *et al.*, 2020. Global shifts in mammalian population trends reveal key predictors of virus spillover risk. *Proceedings of the Royal Society B: biological sciences*, 287 (1924), 20192736. doi:10.1098/rspb.2019.2736
- Kelly, A.H. and Lezaun, J., 2014. Urban mosquitoes, situational publics, and the pursuit of interspecies separation in Dar es Salaam. *American ethnologist*, 41 (2), 368–383. doi:10.1111/amet.12081
- Keune, H., *et al.*, 2017. The need for European OneHealth/EcoHealth networks. *Archives of public health*, 75, 64. doi:10.1186/s13690-017-0232-6
- Kümpel, N.F., *et al.*, 2015. Ebola and bushmeat: myth and reality. *NWFP update*, 5. <http://forestry.fao.org/focus.com/q/1bqqKZHedDwdkXJuzD/wv>
- Kuo, M., Barnes, M., and Jordan, C., 2019. Do experiences with nature promote learning? Converging evidence of a cause-and-effect relationship. *Frontiers in psychology*, 10, 305. doi:10.3389/fpsyg.2019.00305
- Lau, H., *et al.*, 2020. The positive impact of lockdown in Wuhan on containing the COVID-19 outbreak in China. *Journal of travel medicine*, 27 (3), taaa037. doi:10.1093/jtm/taaa037
- Lohiniva, A.L., *et al.*, 2020. Understanding coronavirus disease (COVID-19) risk perceptions among the public to enhance risk communication efforts: a practical approach for outbreaks, Finland, February 2020. *Eurosurveillance*, 25 (13), 2000317. doi:10.2807/1560-7917.ES.2020.25.13.2000317
- Lynteris, C., 2017. Zoonotic diagrams: mastering and unsettling human-animal relations. *Journal of the Royal Anthropological Institute*, 23 (3), 463–485. doi:10.1111/1467-9655.12649
- Maffioli, E.M., 2020. How is the world responding to the 2019 coronavirus disease compared with the 2014 West African Ebola Epidemic? The importance of China as a player in the global economy. *The American journal of tropical medicine and hygiene*, tpm200135. doi:10.4269/ajtmh.20-0135
- Manderson, L. and Levine, S., 2020. COVID-19, risk, fear, and fall-out. *Medical anthropology*, 39, 367–370. doi:10.1080/01459740.2020.1746301
- Martinez-Alvarez, M., *et al.*, 2020. COVID-19 pandemic in west Africa. *The lancet global health*, 8 (5), e631–e632. doi:10.1016/S2214-109X(20)30123-6
- Mills, J.G., *et al.*, 2017. Urban habitat restoration provides a human health benefit through microbiome rewilding: the microbiome rewilding hypothesis. *Restoration ecology*, 25 (6), 866–872. doi:10.1111/rec.12610
- Moss, R., *et al.*, 2016. Model-informed risk assessment and decision making for an emerging infectious disease in the Asia-Pacific region. *PLoS neglected tropical diseases*, 10 (9), e0005018. doi:10.1371/journal.pntd.0005018
- Nagendra, H., *et al.*, 2018. The urban south and the predicament of global sustainability. *Nature sustainability*, 1, 341–349. doi:10.1038/s41893-018-0101-5
- Nisbet, E.K. and Zelenski, J.M., 2011. Underestimating nearby nature: affective forecasting errors obscure the happy path to sustainability. *Psychological science*, 22 (9), 1101–1106. doi:10.1177/0956797611418527
- Nuwagira, E. and Muzoora, C., 2020. Is Sub-Saharan Africa prepared for COVID-19? *Tropical medicine and health*, 48, 18. doi:10.1186/s41182-020-00206-x
- Otmani, A., *et al.*, 2020. Impact of Covid-19 lockdown on PM<sub>10</sub>, SO<sub>2</sub> and NO<sub>2</sub> concentrations in Salé City (Morocco). *Science of the total environment*, 735, 139541. doi:10.1016/j.scitotenv.2020.139541
- Peak, C.M., *et al.*, 2018. Population mobility reductions associated with travel restrictions during the Ebola epidemic in Sierra Leone: use of mobile phone data. *International journal of epidemiology*, 47 (5), 1562–1570. doi:10.1093/ije/dyy095
- Peeri, N.C., *et al.*, 2020. The SARS, MERS and novel coronavirus (COVID-19) epidemics, the newest and biggest global health threats: what lessons have we learned? *International journal of epidemiology*, dyaa033. doi:10.1093/ije/dyaa033
- Plowright, R.K., *et al.*, 2015. Ecological dynamics of emerging bat virus spillover. *Proceedings of the royal society B: biological sciences*, 282 (1798), 20142124. doi:10.1098/rspb.2014.2124
- Pratesi, I., 2020. *Ecosystem destruction and the rise of pandemics - Protect human health by preserving biodiversity*. Rome: WWF Italy.
- Rabinowitz, P.M., *et al.*, 2008. From “us vs. them” to “shared risk”: can animals help link environmental factors to human health? *EcoHealth*, 5, 224–229. doi:10.1007/s10393-008-0170-4
- Rivkin, L.R., *et al.*, 2019. A roadmap for urban evolutionary ecology. *Evolutionary applications*, 12 (3), 384–398. doi:10.1111/eva.12734
- Rohr, J.R., *et al.*, 2019. Emerging human infectious diseases and the links to global food production. *Nature sustainability*, 2 (6), 445–456. doi:10.1038/s41893-019-0293-3
- Roszak, T.E., Gomes, M.E., and Kanner, A.D., 1995. *Ecopsychology: restoring the earth, healing the mind*. San Francisco: Sierra Club Books.

- Rouffaer, L.O., *et al.*, 2017. Effects of urbanization on host-pathogen interactions, using *Yersinia* in house sparrows as a model. *PLoS one*, 12 (12), e0189509. doi:10.1371/journal.pone.0189509
- Rüegg, S.R., Häslar, B., and Zinsstag, J., eds., 2018. *Integrated approaches to health. A handbook for the evaluation of One Health*. Wageningen: Wageningen Academic Publishers.
- Sallis, J.F., *et al.*, 2016. Use of science to guide city planning policy and practice: how to achieve healthy and sustainable future cities. *The lancet*, 388 (10062), 2936–2947. doi:10.1016/S0140-6736(16)30068-X
- Scott, E.A., *et al.*, 2020. A 21st century view of infection control in everyday settings: moving from the germ theory of disease to the microbial theory of health. *American journal of infection control*. doi:10.1016/j.ajic.2020.05.012
- Seltmann, A., *et al.*, 2017. Habitat disturbance results in chronic stress and impaired health status in forest-dwelling paleotropical bats. *Conservation physiology*, 5 (1), cox020. doi:10.1093/conphys/cox020
- Smiley Evans, T., *et al.*, 2020. Synergistic China–US ecological research is essential for global emerging infectious disease preparedness. *EcoHealth*, 17, 160–173. doi:10.1007/s10393-020-01471-2
- Tanne, J.H., *et al.*, 2020. Covid-19: how doctors and health-care systems are tackling coronavirus worldwide. *BMJ*, 368, m1090. doi:10.1136/bmj.m1090
- Thwaites, K., *et al.*, 2005. Restorative urban open space: exploring the spatial configuration of human emotional fulfilment in urban open space. *Landscape research*, 30 (4), 525–547. doi:10.1080/01426390500273346
- Thys, S., 2019a. Contesting the (super)natural origins of Ebola in Macenta, Guinea: biomedical and popular approaches. In: C. Lynteris, ed.. *Framing animals as epidemic villains*. Cham: Palgrave Macmillan, 177–210.
- Thys, S., 2019b. *The added value of anthropology for integrated control of neglected zoonotic diseases in the One Health movement*. Merelbeke: Ghent University Faculty of Veterinary Medicine.
- Trinh, P., *et al.*, 2018. One health relationships between human, animal, and environmental microbiomes: a mini-review. *Frontiers in public health*, 6, 235. doi:10.3389/fpubh.2018.00235
- Vakoch, D.A. and Castrillón, F., eds., 2014. *Ecopsychology, phenomenology, and the environment: the experience of nature*. New York: Springer Science + Business Media. doi:10.1007/978-1-4614-9619-9
- Versini, M., *et al.*, 2015. Unraveling the hygiene hypothesis of helminthes and autoimmunity: origins, pathophysiology, and clinical applications. *BMC medicine*, 13 (1), 81. doi:10.1186/s12916-015-0306-7
- Wang, Q. and Su, M., 2020. A preliminary assessment of the impact of COVID-19 on environment – A case study of China. *Science of the total environment*, 728, 138915. doi:10.1016/j.scitotenv.2020.138915
- Whitburn, J., Linklater, W.L., and Milfont, T.L., 2019. Exposure to urban nature and tree planting are related to pro-environmental behavior via connection to nature, the use of nature for psychological restoration, and environmental attitudes. *Environment and behavior*, 51 (7), 787–810. doi:10.1177/0013916517751009
- Wilson, N., *et al.*, 2020. *Potential Health Impacts from the COVID-19 Pandemic for New Zealand if Eradication Fails: Report to the NZ Ministry of Health*. Available at: [https://www.health.govt.nz/system/files/documents/publications/report\\_for\\_moh\\_-\\_covid-19\\_pandemic\\_nz\\_final.pdf](https://www.health.govt.nz/system/files/documents/publications/report_for_moh_-_covid-19_pandemic_nz_final.pdf)
- World Health Organisation, 2016. *Urban green spaces and health. A review of evidence*. Copenhagen: WHO Regional Office for Europe.
- Zylstra, M.J., *et al.*, 2014. Connectedness as a core conservation concern: an interdisciplinary review of theory and a call for practice. *Springer science reviews*, 2 (1–2), 119–143. doi:10.1007/s40362-014-0021-3